

10 Figures from Introduction and from Experimental Approaches to DNA structure and dynamics




Figure 1:




Figure 2:




Figure 3:

Figure 4:

Figure 5:




Figure 9.10 Electron micrograph of a portion of a 2% agarose gel, 1 mm \times 0.5 mm overall: the small black rectangle is 1000 Å \times 500 Å. Individual gel fibers are about 100 Å wide. Courtesy of Sue Whytock and John Finch.

Figure 6:




Figure 7:




Figure 1.7. Demonstration of the effect of A-tracts in DNA curvature. DNA molecules containing phased A-tract sequences (orange boxes) are cut with a series of restriction enzymes (1–5 in (a)). When analysed by polyacrylamide gel electrophoresis (c), the fragments show varying mobility depending on the position of the A-tract sequences (b).

Figure 8:




Figure 2.6 The significance of phasing in DNA bending. An A tract will introduce a small bend or deflection of the helix axis in DNA. However, the phenomenon of “bent DNA” or “stably curved DNA” that exhibits anomalous migration on polyacrylamide gel electrophoresis requires a number of small individual bends that are in phase. If bends are random, as shown in the top molecule, the DNA will migrate true to its length in an acrylamide gel. Only when bends are phased by 10.5 bp is the stable curvature shown in the middle structure observed. This DNA migrates anomalously slowly on electrophoresis in an acrylamide gel. If, as shown in the bottom structure, bends are phased by 16 bp, successive bends will be directed alternatively up and down. This creates a zig-zag molecule, which is unusually straight and migrates slightly more rapidly than unbent DNA (or DNA containing random bends) in an acrylamide gel.

Figure 9:




Figure 10:




Figure 11:




Figure 12:




Figure 13: